

2003 Health and Preventive Practices Status Report

from Missouri County-level Study



Grundy County**

	County Prevalence (%)*	Northeast Region Prevalence (%)*	State Prevalence (%)*
G eneral			
Health Status (Fair or Poor)	23.4	20.7	16.9
No Health Coverage	13.2	15.4	12.3
B ehavioral Risk Factors			
Physical Inactivity	25.4	25.2	23.0
Current Smoker	27.0	28.2	26.5
Overweight (BMI 25-29.9)	41.0	34.9	35.3
Obese (BMI ≥ 30)	22.7	25.7	23.4
D isease and Conditions			
High Blood Pressure	31.1	29.0	28.5
High Cholesterol (35 years or older)	38.7	39.1	37.3
Asthma	18.6	10.4	10.7
Diabetes	9.7	6.3	7.2
Activity Limitation	19.9	17.8	17.8
C ancer Screening			
Women			
Never had a Mammogram	27.5	17.4	13.4
No Mammogram or Clinical Breast Exam in Last Year:			
40-49 years of age	58.6	59.8	49.4
50-64 years of age	39.4	44.1	35.9
65 and older	51.0	57.5	47.4
Never had a Pap Test:			
18-69 years of age	2.2	5.6	5.1
70 and older	7.9	12.6 H	6.4
No Pap Test in Last Year (18-29)	28.6	23.8	25.8
No Pap Test in Last 3 Years (30-69)	9.3	19.0	13.8
Men and Women 50 Years or Older			
Blood Stool Test:			
Never had a Blood Stool Test	48.4	64.5	57.0
No Blood Stool Test in Last Year	85.5	84.9 H	77.5
Sigmoidoscopy/Colonoscopy			
Never had a Sigmoidoscopy or Colonoscopy	59.8	60.8 H	49.2
No Sigmoidoscopy or Colonoscopy in Past Five Years	65.5	68.5 H	57.0

*Prevalence rates are age adjusted to the 2000 U.S. standard population

■ indicates that the prevalence rate is not significantly above or below state prevalence

H indicates that the prevalence rate is significantly above the state prevalence

L indicates that the prevalence rate is significantly below the state prevalence

Unless otherwise noted respondents are 18 years of age or older

** The best indicators for Grundy county were obtained by combining data from Grundy and Mercer counties.